

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/002, C12N 1/10	A1	(11) International Publication Number: WO 95/25541 (43) International Publication Date: 28 September 1995 (28.09.95)
<p>(21) International Application Number: PCT/US95/03174</p> <p>(22) International Filing Date: 14 March 1995 (14.03.95)</p> <p>(30) Priority Data: 08/215,858 21 March 1994 (21.03.94) US 08/327,516 20 October 1994 (20.10.94) US</p> <p>(71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US).</p> <p>(72) Inventors: CONRAD, Patricia, A.; 3217 Breton Avenue, Davis, CA 95616 (US). BARR, Bradd, C.; 2911 Avila Bay Place, Davis, CA 95616 (US). ANDERSON, Mark, L.; 652 Portsmouth Drive, Davis, CA 95616 (US). SVERLOW, Karen, W.; 249 Wrentham Drive, Vacaville, CA 95688 (US).</p> <p>(74) Agent: BERLINER, Robert; Robbins, Berliner & Carson, 5th floor, 201 N. Figueroa Street, Los Angeles, CA 90012 (US).</p>		<p>(81) Designated States: AU, CA, JP, NZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>

(54) Title: BOVINE NEOSPORA ISOLATES AND THEIR USES

(57) Abstract

The present invention provides isolated bovine *Neospora* cultures. The cultures are used to develop diagnostic assays for the detection of *Neospora* infections in cattle and other animals. Also provided are pharmaceutical compositions for the treatment and prevention of *Neospora* infections.

Reference C

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Bovine Neospora Isolates and Their Uses

This application is a continuation in part of U.S.S.N. 08/215,858, which is incorporated herein by
5 reference.

BACKGROUND OF THE INVENTION

This invention relates to the diagnosis and prevention of bovine diseases caused by the protozoan parasite, *Neospora*. The invention specifically relates
10 to isolated cultures of the parasite and nucleic acids and proteins isolated from them.

A distinct pattern of inflammatory lesions, consisting of focal non-suppurative necrotizing encephalitis, non-suppurative myocarditis and myositis have
15 been observed in many aborted bovine fetuses submitted for diagnosis. The pattern of lesions, particularly in the brain, is similar to those seen with *Toxoplasma gondii* infections in sheep. However, cattle have been reported to be resistant to *T. gondii* infection Dubey,
20 Vet. Parasit. 22:177-202 (1986). In 1988, a cyst-forming protozoal parasite was first identified by histopathological examination in fetuses (Barr et al., Vet. Parasit. 27:354-61 (1990)). This parasite was morphologically similar to *Toxoplasma*, except that some
25 of the cysts had thick walls, which was more similar to the *Neospora caninum*-like protozoan observed by Thilsted & Dubey (*J. Vet. Diagnos. Invest.* 1:205-9 (1989)) in aborted fetuses from a dairy in New Mexico.

Further studies showed that the protozoal
30 parasites associated with inflammatory lesions in aborted fetuses and neonatal calves in California had ultrastructural and antigenic features that were most similar to *N. caninum* parasites which were originally isolated from dogs (Dubey et al., *JAVMA* 193:1259-63
35 (1988)). However, differences in the antigenic reactivity of the bovine protozoan and *N. caninum* when tested with a panel of antisera indicated that they may

not be the same species (Barr et al., Vet. Pathol. 28:110-16 (1991)).

A more complete understanding of the identity and biology of these bovine protozoa requires
5 establishing continuous in vitro cultures of the parasites. Such cultures would also be valuable in the development of diagnostic assays and pharmaceutical compositions for the treatment and prevention of *Neospora* infections. The present invention addresses these and
10 other needs.

SUMMARY OF THE INVENTION

The present invention provides biologically pure cultures of bovine *Neospora*. Two such cultures have
15 been deposited with the ATCC and given ATCC Accession No. _____ (BPA1), and ATCC Accession No. _____ (BPA6).

The invention also provides methods of detecting the presence of antibodies specifically
20 immunoreactive with a bovine *Neospora* antigen in a biological sample (e.g., bovine serum). The method comprises contacting the sample with the *Neospora* antigen, thereby forming an antigen/antibody complex, and detecting the presence or absence of the complex. The
25 *Neospora* antigen is an isolated *Neospora* tachyzoite, bradyzoite, or an isolated *Neospora* protein. In some embodiments, the antigen is immobilized on a solid surface and the complex is detected using a fluorescently labeled anti-bovine antibody.

30 The invention further provides methods of detecting the presence of *Neospora* in a biological sample. These methods comprise contacting the sample with an antibody specifically immunoreactive with a *Neospora* antigen, thereby forming an antigen/antibody
35 complex, and detecting the presence or absence of the complex. The antibody (e.g., a monoclonal antibody) may be immobilized on a solid surface and the complex

detected using a second labeled antibody. Typically, the biological sample is bovine fetal neurological tissue.

The methods of the invention also include detecting the presence of *Neospora*-specific nucleic acids in a biological sample by contacting the sample with a
5 oligonucleotide probe which specifically hybridizes with a target *Neospora*-specific polynucleotide sequence and detecting the presence or absence of hybridization complexes. The methods may further comprise amplifying
10 the target *Neospora*-specific polynucleotide sequence.

The invention further provides pharmaceutical composition comprising a pharmaceutically acceptable carrier and an immunogenically effective amount of a bovine *Neospora* antigen, such as a recombinantly produced
15 bovine *Neospora* polypeptide.

The pharmaceutical compositions are used in protecting a bovine animal from infection by bovine *Neospora*. The compositions are preferably administered to a cow or heifer when the animal is breeding. The
20 pharmaceutical composition is usually administered parenterally.

Definitions

"Antibody" refers to an immunoglobulin molecule able to bind to a specific epitope on an antigen.
25 Antibodies can be a polyclonal mixture or monoclonal. Antibodies can be intact immunoglobulins derived from natural sources or from recombinant sources and can be immunoreactive portions of intact immunoglobulins. Antibodies may exist in a variety of forms including, for
30 example, Fv, F_{ab}, and F(ab)₂, as well as in single chains. Single-chain antibodies, in which genes for a heavy chain and a light chain are combined into a single coding sequence, may also be used.

"Biological sample" refers to any sample
35 obtained from a living or dead organism. Examples of biological samples include biological fluids and tissue specimens. Examples of tissue specimens include fetal

brain tissue, spinal cord, and placenta. Examples of biological fluids include blood, serum, plasma, urine, ascites fluid, cerebrospinal fluid and fetal fluid.

A "biologically pure bovine Neospora culture" refers to a continuous in vitro culture of bovine Neospora organisms (e.g. tachyzoites) which is substantially free of other organisms other than the host cells in which Neospora tachyzoites are grown. A culture is substantially free of other organisms if standard harvesting procedures (as described below) result in a preparation which comprises at least about 95%, preferably 99% or more of the organism, e.g., Neospora tachyzoites.

"Bovine Neospora" refers to Neospora or "Neospora-like" protozoans identified in or isolated from bovine tissues and fluids. Typically, the protozoal parasites can be isolated from neurological tissue of aborted bovine fetuses or congenitally infected calves. Exemplary isolates have been deposited with the American Type Culture Collection, as described below.

A bovine Neospora "protein" or "polypeptide" includes allelic variations normally found in the natural population and changes introduced by recombinant techniques. Those of skill recognize that proteins can be modified in a variety of ways including the addition, deletion and substitution of amino acids.

"Nucleic acids" and "polynucleotides", as used herein, may be DNA or RNA. One of skill will recognize that for use in the expression of Neospora proteins or as diagnostic probes, polynucleotide sequences need not be identical and may be substantially identical to sequences disclosed here. In particular, where the inserted polynucleotide sequence is transcribed and translated to produce a functional polypeptide, one of skill will recognize that because of codon degeneracy a number of polynucleotide sequences will encode the same polypeptide.

"Percentage of sequence identity" for polynucleotides and polypeptides is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Optimal alignment of sequences for comparison may be conducted by computerized implementations of known algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

Substantial identity of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 75% sequence identity, preferably at least 80%, more preferably at least 90% and most preferably at least 95%. Typically, two polypeptides are considered to be substantially identical if at least 40%, preferably at least 60%, more preferably at least 90%, and most preferably at least 95% are identical or conservative substitutions. Sequences are preferably compared to a reference sequence using GAP using default parameters.

Another indication that polynucleotide sequences are substantially identical is if two molecules hybridize to each other under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally,

stringent conditions are selected to be about 5° C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically stringent conditions for a Southern blot protocol involve washing at room temperature with a 5XSSC, 0.1% SDS wash.

The phrase "selectively hybridizing to", refers to a hybridization between a probe and a target sequence in which the probe binds substantially only to the target sequence when the target is in a heterogeneous mixture of polynucleotides and other compounds. Such hybridization is determinative of the presence of the target sequence. Although the probe may bind other unrelated sequences, at least 90%, preferably 95% or more of the hybridization complexes formed are with the target sequence.

The phrase "specifically immunoreactive with", when referring to a protein or peptide, refers to a binding reaction between the protein and an antibody which is determinative of the presence of the protein in the presence of a heterogeneous population of proteins and other compounds. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein and do not bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein and are described in detail below.

The phrase "substantially pure" or "isolated" when referring to a *Neospora* peptide or protein, means a chemical composition which is free of other subcellular components of the *Neospora* organism. Typically, a monomeric protein is substantially pure when at least about 85% or more of a sample exhibits a single

polypeptide backbone. Minor variants or chemical modifications may typically share the same polypeptide sequence. Depending on the purification procedure, purities of 85%, and preferably over 95% pure are possible. Protein purity or homogeneity may be indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualizing a single polypeptide band on a polyacrylamide gel upon silver staining. For certain purposes high resolution will be needed and HPLC or a similar means for purification utilized.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows indirect fluorescent antibody (IFA) titers of serial samples from four cows that aborted *Neospora*-infected fetuses.

Figure 2 shows IFA test titers of serial samples from two cows that aborted *Neospora*-infected fetuses and subsequently delivered congenitally infected calves.

Figure 3 shows seroconversion by heifers experimentally infected with *Neospora*.

Figure 4 shows IFA test titers: comparing cattle with no evidence of infection to dams with *Neospora*-infected fetuses or calves.

Figure 5 shows the nucleotide sequence (SEQ ID No.:1) of DNA encoding nuclear small subunit ribosomal RNA isolated from the cultures of the invention.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention provides *Neospora* cultures isolated from cattle. The cultures are useful in a variety of applications including the production of nucleic acids or proteins for diagnostic assays and the preparation of immunogenic proteins for use in vaccine compositions.

Neospora tachyzoite cultures of the invention have been deposited with the American Type Culture Collection, Rockville, Maryland on March 17, 1994 and given Accession Numbers ____ (BPA1) and ____ (BPA6).

5 ~~These isolates were obtained from tissue~~
homogenates of brain and/or spinal cord of an aborted
bovine fetus and congenitally infected calves.
Immunohistochemistry was used to identify the tachyzoite
and/or cysts associated with lesions in the tissues of
10 these fetuses and calves as Neospora parasites prior to
isolation. Tachyzoite stages of the isolates were grown
in stationary monolayer cultures of bovine fetal
trophoblasts, aortic endothelial cells and/or
macrophages. Electron microscopic studies were used to
15 characterize the ultrastructural features of the BPA1
isolate. Antigenically, tachyzoites of 5 separate
isolates react strongly with antisera to Neospora and
show little or no reactivity with antisera to *Toxoplasma*
gondii or *Hammondia hammondi*. Based on these
20 ultrastructural and antigenic characteristics, these
parasites can be distinguished from the most closely
related and morphologically similar genera of protozoa,
Toxoplasma, *Hammondia* and *Sarcocystis*.

In addition, partial sequences (500-550
25 base pairs) of the 5' end of the nuclear small subunit
(nss)-rRNA gene for three of the bovine Neospora isolates
(BPA1, BPA3 and BPA5) have been obtained and shown to be
identical. The more complete 1.8 kilobase sequence of
the nss-rRNA gene of the BPA1 isolate was obtained and
30 compared to the sequences for this gene in other
coccidial parasites. Alignment of these sequences with
published sequences of *Neospora caninum*, *Cryptosporidium*
parvum, *Sarcocystis muris* and *Toxoplasma gondii* showed
that the bovine Neospora isolate is genotypically unique.

35 As explained in detail below, the isolates are
used to develop a variety of diagnostic assays as well as

pharmaceutical compositions for treatment and prevention of infection.

Preparation of Neospora polypeptides and nucleic acids

5 Standard protein purification techniques can be used to isolate proteins from the tachyzoites or bradyzoites derived from the cultures provided here. Such techniques include selective precipitation with such substances as ammonium sulfate, column chromatography, 10 immunopurification methods, and the like. See, for instance, R. Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag: New York (1982).

 Using standard immunoblot techniques 11 proteins with molecular weights of approximately 106, 15 49.5, 48, 33, 32.5, 30, 28, 26, 19, 18 and 17 kilodaltons (kd) have been identified. All of these proteins are specifically recognized by antibodies from *Neospora* infected cattle. Standard protein purification methods can be used to purify these proteins and produce 20 polyclonal or monoclonal antibodies for use in diagnostic methods described below. Two of these antigens (approximately 106 and 19 kd) have been shown to be useful in enzyme-linked immunoassays (ELISA) for the detection of *Neospora*-specific antibodies in infected 25 cattle.

 Rather than extract the proteins directly from cultured tachyzoites, nucleic acids derived from the cultures can be used for recombinant expression of the proteins. In these methods, the nucleic acids encoding 30 the proteins of interest are introduced into suitable host cells, followed by induction of the cells to produce large amounts of the protein. The invention relies on routine techniques in the field of recombinant genetics, well known to those of ordinary skill in the art. A 35 basic text disclosing the general methods of use in this invention is Sambrook et al., *Molecular Cloning*, A

Laboratory Manual, Cold Spring Harbor Publish., Cold Spring Harbor, NY 2nd ed. (1989).

Nucleic acids for use as diagnostic oligonucleotide probes or for the recombinant expression of proteins can be isolated using a number of techniques. For instance, portions of proteins isolated from the cultures discussed above can be sequenced and used to design degenerate oligonucleotide probes to screen a cDNA library. Amino acid sequencing is performed and oligonucleotide probes are synthesized according to standard techniques as described, for instance, in Sambrook et al., *supra*. Alternatively, oligonucleotide probes useful for identification of desired genes can also be prepared from conserved regions of related genes in other species.

Alternatively, polymerase chain reaction technology (PCR) can be used to amplify nucleic acid sequences of the desired gene directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. Polymerase chain reaction (PCR) or other *in vitro* amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the mRNA in physiological samples, for nucleic acid sequencing, or for other purposes. For a general overview of PCR see PCR Protocols: A Guide to Methods and Applications. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds.), Academic Press, San Diego (1990).

Standard transfection methods are used to produce prokaryotic, mammalian, yeast or insect cell lines which express large quantities of the desired polypeptide, which is then purified using standard techniques. See, e.g., Colley et al., *J. Biol. Chem.* 264:17619-17622, 1989; and Guide to Protein Purification, *supra*.

The nucleotide sequences used to transfect the host cells can be modified to yield *Neospora* polypeptides with a variety of desired properties. For example, the polypeptides can vary from the naturally-occurring
5 sequence at the primary structure level by amino acid, insertions, substitutions, deletions, and the like. These modifications can be used in a number of combinations to produce the final modified protein chain.

The amino acid sequence variants can be
10 prepared with various objectives in mind, including facilitating purification and preparation of the recombinant polypeptide. The modified polypeptides are also useful for modifying plasma half life, improving therapeutic efficacy, and lessening the severity or
15 occurrence of side effects during therapeutic use. The amino acid sequence variants are usually predetermined variants not found in nature but exhibit the same immunogenic activity as naturally occurring protein. In general, modifications of the sequences encoding the
20 polypeptides may be readily accomplished by a variety of well-known techniques, such as site-directed mutagenesis (see, Gillman and Smith, *Gene* 8:81-97, 1979) and Roberts, S. et al., *Nature* 328:731-734, 1987). One of ordinary skill will appreciate that the effect of many mutations
25 is difficult to predict. Thus, most modifications are evaluated by routine screening in a suitable assay for the desired characteristic. For instance, the effect of various modifications on the ability of the polypeptide to elicit a protective immune response can be easily
30 determined using in vitro assays. For instance, the polypeptides can be tested for their ability to induce lymphoproliferation, T cell cytotoxicity, or cytokine production using standard techniques.

The particular procedure used to introduce the
35 genetic material into the host cell for expression of the polypeptide is not particularly critical. Any of the well known procedures for introducing foreign nucleotide

sequences into host cells may be used. These include the use of calcium phosphate transfection, spheroplasts, electroporation, liposomes, microinjection, plasmid vectors, viral vectors and any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell (see Sambrook et al., supra). It is only necessary that the particular procedure utilized be capable of successfully introducing at least one gene into the host cell which is capable of expressing the gene.

Any of a number of well known cells and cell lines can be used to express the polypeptides of the invention. For instance, prokaryotic cells such as *E. coli* can be used. Eukaryotic cells include, yeast, Chinese hamster ovary (CHO) cells, COS cells, and insect cells.

The particular vector used to transport the genetic information into the cell is also not particularly critical. Any of the conventional vectors used for expression of recombinant proteins in prokaryotic and eukaryotic cells may be used. Expression vectors for mammalian cells typically contain regulatory elements from eukaryotic viruses.

The expression vector typically contains a transcription unit or expression cassette that contains all the elements required for the expression of the polypeptide DNA in the host cells. A typical expression cassette contains a promoter operably linked to the DNA sequence encoding a polypeptide and signals required for efficient polyadenylation of the transcript. The term "operably linked" as used herein refers to linkage of a promoter upstream from a DNA sequence such that the promoter mediates transcription of the DNA sequence. The promoter is preferably positioned about the same distance from the heterologous transcription start site as it is from the transcription start site in its natural setting.

As is known in the art, however, some variation in this distance can be accommodated without loss of promoter function.

Following the growth of the recombinant cells and expression of the polypeptide, the culture medium is harvested for purification of the secreted protein. The media are typically clarified by centrifugation or filtration to remove cells and cell debris and the proteins are concentrated by adsorption to any suitable resin or by use of ammonium sulfate fractionation, polyethylene glycol precipitation, or by ultrafiltration. Other routine means known in the art may be equally suitable. Further purification of the polypeptide can be accomplished by standard techniques, for example, affinity chromatography, ion exchange chromatography, sizing chromatography, His₆ tagging and Ni-agarose chromatography (as described in Dobeli et al. *Mol. and Biochem. Parasit.* 41:259-268 (1990)), or other protein purification techniques to obtain homogeneity. The purified proteins are then used to produce pharmaceutical compositions, as described below.

An alternative method of preparing recombinant polypeptides useful as vaccines involves the use of recombinant viruses (e.g., vaccinia). Vaccinia virus is grown in suitable cultured mammalian cells such as the HeLA S3 spinner cells, as described by Mackett, Smith and Moss, "The construction and characterization of Vaccinia Virus Recombinants Expressing Foreign Genes" in "DNA cloning Vol. II. A practical approach", Ed. D.M. Glover, IRL Press, Oxford, pp 191-211.

Antibody Production

The isolated proteins or cultures of the present invention can be used to produce antibodies specifically reactive with *Neospora* antigens. If isolated proteins are used, they may be recombinantly produced or isolated from *Neospora* cultures. Synthetic

peptides made using the protein sequences may also be used.

Methods of production of polyclonal antibodies are known to those of skill in the art. In brief, an immunogen, preferably a purified protein, is mixed with an adjuvant and animals are immunized. When appropriately high titers of antibody to the immunogen are obtained, blood is collected from the animal and antisera is prepared. Further fractionation of the antisera to enrich for antibodies reactive to *Neospora* proteins can be done if desired. (See Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Pubs., N.Y. (1988)).

Polyclonal antisera to the BPA1 and BPA3 isolates have been produced and evaluated. The polyclonal antisera are used to identify and characterize *Neospora* tachyzoite and bradyzoite stages in the tissues of infected animals using, for instance, immunoperoxidase test procedures described in Anderson et al. *JAVMA* 198:241 (1991) and Barr et al. *Vet. Pathol.* 28:110-116(1991).

Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell (See, Kohler and Milstein, *Eur. J. Immunol.* 6:511-519 (1976)). Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host.

For instance, the BPA1 isolate has been used to immunize mice to obtain sensitized B cells for hybridoma production. Using these cells, monoclonal antibodies to the 48 kd and 70 kd *Neospora* proteins have been obtained. The monoclonal antibodies produced are used, for instance, in ELISA diagnostic tests, immunohistochemical tests, for the *in vitro* evaluation of parasite invasion, to select candidate antigens for vaccine development, protein isolation, and for screening genomic and cDNA libraries to select appropriate gene sequences.

Diagnosis of *Neospora* infections

The present invention also provides methods for detecting the presence or absence of *Neospora* in a biological sample. For instance, antibodies specifically reactive with *Neospora* can be detected using either proteins or the isolates described here. The proteins and isolates can also be used to raise specific antibodies (either monoclonal or polyclonal) to detect the antigen in a sample. In addition, the nucleic acids disclosed and claimed here can be used to detect *Neospora*-specific sequences using standard hybridization techniques. Each of these assays is described below.

A. Immunoassays

For a review of immunological and immunoassay procedures in general, see *Basic and Clinical Immunology* 7th Edition (D. Stites and A. Terr ed.) 1991. The immunoassays of the present invention can be performed in any of several configurations, which are reviewed extensively in *Enzyme Immunoassay*, E.T. Maggio, ed., CRC Press, Boca Raton, Florida (1980); "Practice and Theory of Enzyme Immunoassays," P. Tijssen, *Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers B.V. Amsterdam (1985). For instance, the proteins and antibodies disclosed here are conveniently used in ELISA, immunoblot analysis and agglutination assays. Particularly preferred assay

formats include the indirect fluorescent antibody assay as described in Example 2.

In brief, immunoassays to measure anti-Neospora antibodies or antigens can be either competitive or noncompetitive binding assays. In competitive binding assays, the sample analyte (e.g., anti-Neospora antibodies) competes with a labeled analyte (e.g., anti-Neospora monoclonal antibody) for specific binding sites on a capture agent (e.g., isolated Neospora protein) bound to a solid surface. The concentration of labeled analyte bound to the capture agent is inversely proportional to the amount of free analyte present in the sample.

Noncompetitive assays are typically sandwich assays, in which the sample analyte is bound between two analyte-specific binding reagents. One of the binding agents is used as a capture agent and is bound to a solid surface. The second binding agent is labelled and is used to measure or detect the resultant complex by visual or instrument means.

A number of combinations of capture agent and labelled binding agent can be used. For instance, an isolated Neospora protein or culture can be used as the capture agent and labelled anti-bovine antibodies specific for the constant region of bovine antibodies can be used as the labelled binding agent. Goat, sheep and other non-bovine antibodies specific for bovine immunoglobulin constant regions (eg. γ or μ) are well known in the art. Alternatively, the anti-bovine antibodies can be the capture agent and the antigen can be labelled.

Various components of the assay, including the antigen, anti-Neospora antibody, or anti-bovine antibody, may be bound to a solid surface. Many methods for immobilizing biomolecules to a variety of solid surfaces are known in the art. For instance, the solid surface may be a membrane (e.g., nitrocellulose), a microtiter dish (e.g., PVC or polystyrene) or a bead. The desired

component may be covalently bound or noncovalently attached through nonspecific bonding.

Alternatively, the immunoassay may be carried out in liquid phase and a variety of separation methods may be employed to separate the bound labeled component from the unbound labelled components. These methods are known to those of skill in the art and include immunoprecipitation, column chromatography, adsorption, addition of magnetizable particles coated with a binding agent and other similar procedures.

An immunoassay may also be carried out in liquid phase without a separation procedure. Various homogeneous immunoassay methods are now being applied to immunoassays for protein analytes. In these methods, the binding of the binding agent to the analyte causes a change in the signal emitted by the label, so that binding may be measured without separating the bound from the unbound labelled component.

Western blot (immunoblot) analysis can also be used to detect the presence of antibodies to *Neospora* in the sample. This technique is a reliable method for confirming the presence of antibodies against a particular protein in the sample. The technique generally comprises separating proteins by gel electrophoresis on the basis of molecular weight, transferring the separated proteins to a suitable solid support, (such as a nitrocellulose filter, a nylon filter, or derivatized nylon filter), and incubating the sample with the separated proteins. This causes specific target antibodies present in the sample to bind their respective proteins. Target antibodies are then detected using labeled anti-bovine antibodies.

The immunoassay formats described above employ labelled assay components. The label can be in a variety of forms. The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. A wide

variety of labels may be used. The component may be labelled by any one of several methods. Traditionally a radioactive label incorporating ^3H , ^{125}I , ^{35}S , ^{14}C , or ^{32}P was used. Non-radioactive labels include ligands which
5 bind to labelled antibodies, fluorophores, chemiluminescent agents, enzymes, and antibodies which can serve as specific binding pair members for a labelled ligand. The choice of label depends on sensitivity required, ease of conjugation with the compound,
10 stability requirements, and available instrumentation.

Enzymes of interest as labels will primarily be hydrolases, particularly phosphatases, esterases and glycosidases, or oxidoreductases, particularly peroxidases. Fluorescent compounds include fluorescein
15 and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, etc. Chemiluminescent compounds include luciferin, and 2,3-dihydrophthalazinediones, e.g., luminol. For a review of various labelling or signal producing systems which may be used, see U.S.
20 Patent No. 4,391,904, which is incorporated herein by reference.

Non-radioactive labels are often attached by indirect means. Generally, a ligand molecule (e.g., biotin) is covalently bound to the molecule. The ligand
25 then binds to an anti-ligand (e.g., streptavidin) molecule which is either inherently detectable or covalently bound to a signal system, such as a detectable enzyme, a fluorescent compound, or a chemiluminescent compound. A number of ligands and anti-ligands can be
30 used. Where a ligand has a natural anti-ligand, for example, biotin, thyroxine, and cortisol, it can be used in conjunction with the labelled, naturally occurring anti-ligands. Alternatively, any haptenic or antigenic compound can be used in combination with an antibody.

35 Some assay formats do not require the use of labelled components. For instance, agglutination assays can be used to detect the presence of the target

antibodies. In this case, antigen-coated particles are agglutinated by samples comprising the target antibodies. In this format, none of the components need be labelled and the presence of the target antibody is detected by
5 simple visual inspection.

B. Detection of Neospora nucleic acids

As noted above, this invention also embraces methods for detecting the presence of Neospora DNA or RNA in biological samples. These sequences can be used to
10 detect all stages of the Neospora life cycle (e.g., tachyzoites, bradyzoites, and oocysts) in biological samples from both the bovine host and the definitive host. A variety of methods of specific DNA and RNA measurement using nucleic acid hybridization techniques
15 are known to those of skill in the art. See Sambrook et al., supra.

One method for determining the presence or absence of Neospora DNA in a sample involves a Southern transfer. Briefly, the digested DNA is run on agarose
20 slab gels in buffer and transferred to membranes. In a similar manner, a Northern transfer may be used for the detection of Neospora mRNA in samples of RNA. Hybridization is carried out using labelled oligonucleotide probes which specifically hybridize to
25 Neospora nucleic acids. Labels used for this purpose are generally as described for immunoassays. Visualization of the hybridized portions allows the qualitative determination of the presence or absence of Neospora genes.

30 A variety of other nucleic acid hybridization formats are known to those skilled in the art. For example, common formats include sandwich assays and competition or displacement assays. Hybridization techniques are generally described in "Nucleic Acid
35 Hybridization, A Practical Approach," Ed. Hades, B.D. and Higgins, S.J., IRL Press, 1985; Gall and Pardue (1969),

Proc. Natl. Acad. Sci., U.S.A., 63:378-383; and John, Burnsteil and Jones (1969) *Nature*, 223:582-587.

Sandwich assays are commercially useful hybridization assays for detecting or isolating nucleic acid sequences. Such assays utilize a "capture" nucleic acid covalently immobilized to a solid support and labelled "signal" nucleic acid in solution. The biological sample will provide the target nucleic acid. The "capture" nucleic acid and "signal" nucleic acid probe hybridize with the target nucleic acid to form a "sandwich" hybridization complex. To be effective, the signal nucleic acid cannot hybridize with the capture nucleic acid.

The sensitivity of the hybridization assays may be enhanced through use of a nucleic acid amplification system which multiplies the target nucleic acid being detected. Examples of such systems include the polymerase chain reaction (PCR) system and the ligase chain reaction (LCR) system. Other methods recently described in the art are the nucleic acid sequence based amplification (NASBA™, Cangene, Mississauga, Ontario) and Q Beta Replicase systems.

An alternative means for detecting *Neospora* nucleic acids is *in situ* hybridization. *In situ* hybridization assays are well known and are generally described in Angerer, et al., *Methods Enzymol.*, 152:649-660 (1987). *In situ* hybridization assays use cells or tissue fixed to a solid support, typically a glass slide. If DNA is to be probed, the cells are denatured with heat or alkali. The cells are then contacted with a hybridization solution at a moderate temperature to permit annealing of labelled *Neospora* specific probes. The probes are preferably labelled with radioisotopes or fluorescent reporters.

Exemplary nucleic acid sequences for use in the assays described above include sequences from the nss-rRNA sequences disclosed here. For instance, the primer

and probe sequences disclosed in Example 4 can be used to amplify and identify nucleic acids of bovine *Neospora* in blood, cerebrospinal fluid and fetal fluids, as well as in frozen or formalin-fixed tissue. These primers are particularly useful for the diagnosis of neosporosis and identification of the source of *Neospora* parasite stages (tachyzoites, bradyzoites and oocysts) in various animal hosts.

Pharmaceutical Compositions comprising *Neospora*

10 A pharmaceutical composition prepared using anti-*Neospora* monoclonal antibodies or fragments thereof as well as *Neospora* proteins or their immunogenic equivalents can be used in a variety of pharmaceutical preparations for the treatment and/or prevention of
15 *Neospora* infections.

The pharmaceutical compositions are typically used to vaccinate cattle, sheep, goats and other animals infected by *Neospora*. The compositions of the invention can also be used to treat the definitive host to prevent
20 the shedding of oocysts and subsequent transfer to cattle. The compositions for administration to either cattle or the definitive host can comprise tachyzoite and/or bradyzoite antigens.

An attenuated *Neospora* vaccine can only be used
25 in the absence of a risk of human infection should the milk or tissues of immunized animals be consumed. Thus, preferred vaccines are subunit vaccines that elicit antibody and cell-mediated immunity (CMI) to antigens of bovine *Neospora*. Experimental evidence indicates that
30 CMI is an important component of the protective immune response in cattle. A variety of methods for evaluating the specificity of the helper and cytotoxic T cell response to selected antigens in vitro can be used. In addition, as demonstrated below, cows infected using
35 culture-derived tachyzoites mount a protective immune response and prevent transplacental infection of the fetus.

The vaccines of the invention are typically administered orally or parenterally, usually intramuscularly or subcutaneously. For parenteral administration, the antigen may be combined with a suitable carrier. For example, it may be administered in water, saline or buffered vehicles with or without various adjuvants or immunomodulating agents such as aluminum hydroxide, aluminum phosphate, aluminum potassium sulfate (alum), beryllium sulfate, silica, kaolin, carbon, water-in-oil emulsions, oil-in-water emulsions, muramyl dipeptide, bacterial endotoxin, lipid, *Bordetella pertussis*, and the like. Such adjuvants are available commercially from various sources, for example, Merck Adjuvant 6 (Merck and Company, Inc., Rahway, N.J.). Other suitable adjuvants are MPL+TDM Emulsion (RIBBI Immunochem Research Inc. U.S.A.). Other immunostimulants include interleukin 1, interleukin 2 and interferon-gamma. These proteins can be provided with the vaccine or their corresponding genetic sequence provided as a functional operon with a recombinant vaccine system such as vaccinia virus. The proportion of antigen and adjuvant can be varied over a broad range so long as both are present in effective amounts.

Vaccine compositions of the invention are administered to a cattle, sheep, or goats susceptible to or otherwise at risk of infection to elicit an immune response against the antigen and thus enhance the patient's own immune response capabilities. Such an amount is defined to be an "immunogenically effective amount." In this use, the precise amounts depend on the judgement of the prescribing veterinarian and would include consideration of the patient's state of health and weight, the mode of administration, the nature of the formulation, and the like. Generally, on a per-dose basis, the concentration of the *Neospora* antigen can range from about 1 μ g to about 100 mg per bovine host. For administration to cattle, a preferable range is from

about 100 μ g to about 1 mg per unit dose. A suitable dose size is about 1-10 ml, preferably about 1.0 ml. Accordingly, a typical dose for subcutaneous injection, for example, would comprise 1 to 2 ml containing 0.1 to 5 10 mg of antigen.

A variety of vaccination regimens may be effective in immunizing cattle and other animals. Preferably, female cattle (heifers and cows) are vaccinated just prior to or at the time of breeding so as 10 to prevent abortion and reduce the possibility of congenital infections. A second immunization will be given 2-4 weeks after initial immunization. Calves and adult males may also be vaccinated, if desired. Animals that have been previously exposed to *Neospora* or have 15 received colostrum antibodies from the mother may require booster injections. The booster injection is preferably timed to coincide with times of maximal challenge and/or risk of abortion. Different immunization regimes may be adopted depending on the judgement of the veterinarian.

20 Vaccines of the invention may comprise a crude extract of *Neospora* tachyzoites, bradyzoites or other stages. Chemically fixed parasites or cells can also be used. As noted above, preferred vaccines comprise partially or completely purified *Neospora* 25 protein preparations. The antigen produced by recombinant DNA technology is preferred because it is more economical than the other sources and is more readily purified in large quantities.

In addition to use in recombinant expression 30 systems, the isolated *Neospora* gene sequences can also be used to transform viruses that transfect host cells in animals. Live attenuated viruses, such as vaccinia or adenovirus, are convenient alternatives for vaccines because they are inexpensive to produce and are easily 35 transported and administered.

Suitable viruses for use in the present invention include, but are not limited to, pox viruses,

such as capripox and cowpox viruses, and vaccinia viruses, alpha viruses, adenoviruses, and other animal viruses. The recombinant viruses can be produced by methods well known in the art, for example, using
5 homologous recombination or ligating two plasmids. A recombinant canarypox or cowpox virus can be made, for example, by inserting the DNA's encoding the Neospora protein or fragments thereof into plasmids so that they are flanked by viral sequences on both sides. The DNA's
10 encoding Neospora polypeptides are then inserted into the virus genome through homologous recombination.

Preferentially, a viral vaccine using recombinant vaccinia virus is used. A vaccine prepared utilizing the gene encoding the Neospora protein
15 incorporated into vaccinia virus would comprise stocks of recombinant virus where the gene encoding the Neospora protein is integrated into the genome of the virus in a form suitable for expression of the gene.

20

EXAMPLES

Example 1

This example describes the isolation and in vitro cultivation of Neospora from aborted bovine
25 fetuses. The isolation of these 2 cultures (BPA1 and BPA2) is described in Conrad et al. *Parasitol.* 106:239-249 (1993). Additional cultures (BPA3-6) were also isolated using the same technique, except that 1 ml (instead of 2 ml) of brain or spinal cord homogenate was
30 trypsinized and then incubated for 2-4 hours, rather than overnight, on the cell monolayer. In addition, the bovine aortic endothelial cell line (CPAE: American Tissue Culture Collection #CCL209) was found to be the best cell monolayer for the cultivation of bovine
35 Neospora. One of these cultures (BPA6) has been shown to induce bradyzoite cysts in mice.

MATERIALS AND METHODS

Pathological examination and immunohistochemistry of fetal tissues

Aborted bovine fetuses submitted to the California Veterinary Diagnostic Laboratory System were necropsied using standard techniques. The brains from fetuses suspected of having protozoal infections were removed aseptically from the cranium. One half of the brain was placed in sterile saline (0-85% w/v) containing 1000 U/ml penicillin G and 100 µg/ml streptomycin (antibiotic saline) and stored at 4 °C until a diagnosis of protozoal infection was confirmed, at which time the brain could be processed for in vitro cultivation. Multiple tissues, including portions of the brain, liver, kidney, heart, lung, spleen, gastrointestinal tract, skeletal muscle, adrenal, trachea and thymus, were collected from each fetus and fixed in 10% neutral buffered formalin for 24 h. Fixed tissues were trimmed, embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined by light microscopy for the presence of lesions and parasites, as previously described (Barr et al. 1990 Vet. Pathol. 27:354-61).

Fetuses with multifocal microgliosis and/or necrosis in the brain, suggesting protozoal infection, were further examined by immunohistochemistry for the presence of parasites in brain tissue sections using an avidin-biotin peroxidase complex procedure (Vector Laboratories, Burlingame, CA, USA) with anti-rabbit serum to detect tissue-binding of rabbit polyclonal anti-N. caninum serum. The immunoperoxidase method employed was basically as described previously (Barr et al. 1991 J. Vet. Diag. Invest. 3:39-46) except that tissue sections were processed using a microprobe system (FisherBiotech, Pittsburgh, PA, USA) and Probe-On glass slides (FisherBiotech). Aminoethylcarbazole (A.E.C. Substrate System, Dako, Santa Barbara, CA, USA) was the chromogen.

Parasites in the tissue sections of brains from the 66th and 93rd fetus (hereafter referred to as fetus 66 and 93) were further characterized by the same immunohistochemical procedure to test their reactivity with antisera to additional apicomplexan protozoal parasites. Tissue sections were incubated at room temperature for 1 h with optimal dilutions of the following antisera: 1:1000 dilution of antiserum to *N. caninum* tachyzoites (Lindsay & Dubey, 1989 Am. J. Vet. Res. 50:1981-3), 1:50 dilution of antiserum to *Hammondia hammondi* tissue cysts and 4 different antisera to *T. gondii* (Tg1-4). Antiserum Tg1 was produced by the infection of a rabbit with live sporulated oocysts of the ME-49 strain (Lindsay & Dubey, supra) of *T. gondii* and used at a 1:400 dilution. *Toxoplasma gondii* antiserum Tg2 (Dr J. C. Boothroyd, Stanford University) was produced by immunization of a rabbit with a tachyzoite lysate of the RH strain of *T. gondii* and was used at a dilution of 1:300. Antisera Tg3 (BioGenex Laboratories, Dublin, CA, USA) and Tg4 (I.C.N. Immunobiologicals, Lisle, IL, USA) were developed by immunizing rabbits with tachyzoites of the RH and H44 strains, respectively. Antiserum Tg3 was applied as supplied by the manufacturer and Tg4 was used at a 1:80 dilution. The optimal dilution chosen for each antiserum produced a strongly positive reaction against the respective positive control parasite with no appreciable non-specific, background staining. Control tissues consisted of paraffin-embedded sections of murine brain with *N. caninum* tachyzoites, murine brain with *T. gondii* cysts, murine spleen with *T. gondii* tachyzoites, murine skeletal muscle with *H. hammondi* cysts and bovine tongue with *Sarcocystis cruzi* cysts (Barr et al. 1991 Vet. Path. 28:110-116).

Parasite cultures

Stationary monolayer cultures of bovine cardiopulmonary aortic endothelial cells (CPAE: ATCC #CCL209) and M617 bovine macrophages (Speer et al. 1985

Infect. and Immun. 50:566-71) were maintained in Dulbecco's Minimum Essential Medium (DMEM:GIBCO Laboratories, Grand Island, NY, USA) supplemented with 10% (v/v) heat-inactivated fetal bovine serum (FBS) or 5 heat-inactivated adult equine serum (HS), 2 mM L-glutamine, 50 U/ml penicillin and 50 µg/ml streptomycin (DMEM-FBS or DMEM-HS). Bovine fetal trophoblast cells (87-3) were grown as previously described by Munson et al. 1988 *J. Tissue Cult. Methods* 11:123-8) in DMEM 10 supplemented with 10% (v/v) FBS, 5 µg/ml transferrin, 5 µg/ml insulin, 5 ng/ml selenium, 10 ng/ml epidermal growth factor, 100 µg/ml streptomycin, 100 U/ml penicillin and 0.25 µg/ml amphotericin B (DMEM-FBS*). Control cultures of *T. gondii* (RH strain provided by Dr 15 J. Boothroyd) and *N. caninum* (NCl; Dubey et al. 1988 *J Am Vet Med Assoc* 193:1259-1263) were maintained in the CPAE and M617 monolayer cultures. Parasite-infected and uninfected monolayer cultures were maintained in 25 or 75 cm² flasks incubated at 37 °C with 5% CO₂. Culture medium 20 was changed 3 times weekly. Established parasite cultures were passaged to uninfected monolayers when 70-90% of cells were infected. To passage parasites, the infected monolayer was removed from the flask by scraping into the media and passed 3 times through a 25-gauge 25 needle to disrupt the cells. The cell suspension was then diluted from 1:2 to 1:8 in fresh complete media and added to a confluent, uninfected monolayer.

After protozoal infection was confirmed by histology, fetal brain tissue that was stored at 4 °C for 30 a variable period of time in antibiotic saline was processed for in vitro cultivation. In all cases where isolates were obtained in vitro, tissue cysts were seen in tissue sections of bovine fetal brain. Half of the fetal brain in approximately 25 ml of antibiotic saline 35 was ground with a pestle and mortar and filtered through sterile gauze. Aliquots of 2 ml of brain homogenate were placed in 10 ml of 0.05% (v/v) trypsin and incubated at

37 °C for 1 h. After trypsin digestion, the material was pelleted by centrifugation at 600 g for 10 min. The supernatant was discarded and the pellet was resuspended in 1-3 ml of either DMEM-HS or DMEM-FBS. Brain from fetus 66 was prepared for culture 48 h after submission and 1 ml of trypsinized brain suspension was dispensed into a 25 cm² flask of bovine 87-3 trophoblast cells. Brain from fetus 93 was processed 10 days after submission when half of the trypsinized brain was placed in a 25 cm² flask of 87-3 trophoblast cells and the remainder in a 75 cm² flask of endothelial cells. After incubation overnight, the brain suspension from both fetuses was removed from the flask and the monolayers were washed 3 times with the appropriate media before adding 5-10 ml of fresh media. Cultures were maintained as described above and examined with an inverted microscope for the presence of parasites.

Immunohistochemistry of tachyzoites in vitro

Antigenic reactivity of the two in vitro isolates from aborted bovine fetuses was compared to that of tachyzoites from control cultures of *T. gondii* and *N. caninum*. Tachyzoites of each isolate were harvested during logarithmic growth by scraping the infected CPAE monolayer from a 25 cm² tissue culture flask. Monolayer cells were disrupted by repeated passage through a 25-gauge needle. The suspension was passed through a 5 µm filter to remove cellular debris and pelleted by centrifugation at 1500 g for 10 min. After removing the supernatant fraction, the pelleted tachyzoites of each isolate were resuspended in DMEM-HS and inoculated into each of the wells on two 4-chambered tissue culture slides (Lab-Tek, Nunc, Naperville, IL, USA). Each of the 4 chambers on these slides were seeded 24-48 h prior to parasite inoculation with CPAE cells so that the monolayers were 60-80% confluent at the time of infection. The appropriate slides were inoculated with the slower growing bovine fetal isolates first to allow

the parasites to grow for 48 h, whereas the isolates of *T. gondii* and *N. caninum* were cultivated on slides for 24 h before being processed for immunohistochemical evaluation.

5 To prepare the parasite cultures for immunohistochemistry, culture supernatants were removed with monolayers remaining adherent to glass slides. These slides were fixed in 100% methanol (4 °C) for 10 min and allowed to air dry completely before being washed
10 3 times for 5 min each in physiologically buffered saline (PBS:pH 7-2), incubated for 10 min in 3% (v/v) hydrogen peroxide in methanol, washed again 3 times for 5 min each in PBS and incubated for 30 min with 20% goat serum to block non-specific antibody binding sites. Each slide
15 was then incubated for 1 h with 3 wells containing different primary antiserum and 1 well serving as a negative control with a pre-infection rabbit serum. The optimal antisera dilutions were selected to produce a strongly positive reaction against the homologous
20 culture-derived antigen with no appreciable non-specific, background staining. The dilutions of antisera used for staining parasites in vitro were 1:3000 for *N. caninum*, 1: 800 for Tg1, 1:40 for Tg2, 1:1 for Tg3, 1:2000 for Tg4 and 1:50 for *H. hammondi*. Slides were washed 3 times for
25 5 min each in PBS and the secondary antibody and conjugate were applied as described above for the tissue sections except that the slides were processed manually and the chromogen was applied for only 2 min.

30 RESULTS

The first parasite isolate (BPA1) was obtained from fetus 66 which was estimated to be approximately 4 months gestational age and in relatively good postmortem condition at the time of necropsy. Significant gross
35 lesions were restricted to focal epicardial petechiae. On histological examination there were infrequent, random, small foci of gliosis and 5 protozoal cysts were

seen in sections of the fetal brain. The tissue cysts ranged from 8 to 10 μm in diameter and had distinct, thin walls ($< 1 \mu\text{m}$) surrounding at least 25-40 closely packed bradyzoites. In addition there were scattered
5 mononuclear inflammatory cell infiltrates in the heart, portal tracts of the liver and throughout the renal cortex. In the lung, macrophages and neutrophils were present within alveolar septa, adjacent to bronchioles and free in the lumen of bronchioles and alveoli.
10 *Escherichia coli* and *Proteus* spp. were isolated from the lung, liver, spleen and abomasal contents of this fetus.

The second isolate (BPA2) was obtained from fetus 93 which had an estimated gestational age of 6 months and was mildly autolyzed. Histological
15 examination revealed infrequent random foci of gliosis in the brain with adjacent capillaries that had prominent hypertrophied endothelium. There was also a mild diffuse meningeal infiltrate of mononuclear cells, consisting predominantly of lymphocytes with occasional plasma
20 cells. Four randomly located protozoal tissue cysts were found in the brain; one being located adjacent to a focus of gliosis. The cysts were 8-13 μm in diameter with at least 25-50 bradyzoites. Two of the cysts had thick (1-2 μm) walls. Focal, mixed mononuclear inflammatory cell
25 infiltrates were also present in skeletal muscle and in the renal cortex.

Table 1 summarizes the immunoreactivity of protozoal tissue cysts in the brains of fetus 66 and fetus 93 with different polyclonal antisera and compares
30 these antigenic reactions to those of *N. caninum*, *T. gondii*, *H. hammondi* and *S. cruzi*. The protozoal cysts in fetuses 66 and 93 reacted most strongly to *N. caninum* antiserum and had weaker reactions to *H. hammondi* antiserum. With both antisera, staining was
35 predominantly to the cyst wall with some staining of zoites within the cysts. Overall, the reactivity of cysts in the two bovine fetuses was more similar to the

reactivity of *N. caninum* tachyzoites than to that of *T. gondii* tachyzoites or cysts, *H. hammondia* cysts or *S. cruzi* cysts.

Table 1 Reactivity of tissue cysts and tachyzoites with rabbit polyclonal antisera against different parasites in an immunoperoxidase test

Parasite antigen	Tissue	Antisera					
		Neospora caninum	Hammondia hammondi	Toxoplasma gondii			
				Tg1	Tg2	Tg3	Tg4
Fetus 66 cysts	Bovine brain	++ [*]	++ [*]	±	-	-	-
Fetus 93 cysts	Bovine brain	++ [*]	+ [*]	-	-	-	-
Neospora caninum tachyzoites	Mouse brain	+++	-	++	-	-	-
Toxoplasma gondii tachyzoites	Mouse spleen	-	++	+++	+++	±	±
Toxoplasma gondii cysts	Mouse brain	+	+++	+++	+	++	++
Hammondia hammondi cysts	Mouse muscle	±	+++	+++	-	+	++
Sarcocystis cruzi cysts	Bovine muscle	-	-	-	-	-	-
*Primarily cyst wall staining.							

In approximately 14 months, over 100 fetuses were submitted specifically as protozoal abortion suspects from dairies in California. Parasite isolations were attempted with brain from the 49 fetuses which had Neospora-like protozoa identified by immunohistochemistry. The first evidence of parasite growth in these cultures was detected in the 87-3 cell line 34 days after inoculation of brain tissue from fetus 66 (isolate BAP1). The next successful in vitro isolation was apparent when tachyzoites were first observed in 87-3 and CPAE cultures on day 15 after inoculation with brain tissue from fetus 93 (isolate BPA2). In cultures of both isolates, parasites first appeared as small clusters of intracellular tachyzoites in pairs or random groups. Extracellular tachyzoites were seen escaping from the bovine cell monolayers and moving by gliding and twisting in the culture medium. On Giemsa-stained smears of infected monolayers,

extracellular tachyzoites were 1.5-2.5 μm wide at the nucleus and 6-8 μm long. The number of tachyzoite clusters and number of tachyzoites in each cluster increased gradually in the cultures as they became established with continuous parasite growth. Generally, 5 parasite clusters contained approximately 10-100 tachyzoites. Growth of both isolates was maintained in cultures of 87-3, CPAE, or M617. However, the best growth was observed in the 87-3 and CPAE cultures. 10 Within 2-3 months of establishment, the cultures were passaged weekly whenever approximately 80-95% of the bovine monolayer cells were infected with tachyzoites. Routinely, the established BPA1 and BPA2 cultures were passaged by adding a 1:8 dilution of needle-passaged 15 monolayer in fresh media to cultures of uninfected bovine monolayers. By comparison, in our laboratory, cultures of *T. gondii* (RH isolate) are routinely passaged at a 1:200 dilution and *N. caninum* (NC-1 isolate) cultures are passaged at a 1:10 dilution every 2-3 days. As of 20 mid-May 1992, cultures of the BPA1 and BPA2 isolates had been maintained with continuous growth for 10 and 6 months, respectively.

The results of an antigenic comparison of in vitro cultivated tachyzoites of BPA1 and BPA2 to those of 25 cultivated *N. caninum* and *T. gondii* tachyzoites are shown in Table 2. The reactions of the bovine fetal isolates to the different antisera were similar to that demonstrated by *N. caninum*, and distinctly different from the pattern of reactivity observed with *T. gondii* 30 tachyzoites (Table 2).

Table 2. Reactivity of in vitro cultivated tachyzoites with rabbit polyclonal antisera against different parasites in an immunoperoxidase test

5	Parasite antigen	Antisera					
		Neospora caninum	Hammondia hammondi	Toxoplasma gondii			
				Tg1	Tg2	Tg3	Tg4
	Foetus 66 (BPA1)	+++	-	-	+	-	-
	Foetus 93 (BPA2)	+++	-	±	+	-	-
10	Neospora caninum	+++	-	-	+	-	-
	Toxoplasma gondii	-	+++	+++	+++	+++	+++

By transmission electron microscopy, the in vitro tachyzoites of isolates BPA1 and BPA2 were morphologically similar. Therefore, the following ultrastructural description applies to both isolates. Individual tachyzoites or clusters of multiple tachyzoites were usually located within a parasitophorous vacuole in the cytoplasm of bovine monolayer cells. Tachyzoites had a pellicle consisting of a complex of 2 inner membranes beneath a plasmalemmal membrane, a prominent nucleus in the central or posterior portion of the tachyzoite, 1 to 3 profiles of long tubular cristate mitochondria, a Golgi complex, rough and smooth endoplasmic reticulum, single- or multiple-membraned vesicles, and numerous free ribosomes. Ultrastructural apical features characteristic of apicomplexan parasites were present in tachyzoites of both isolates, including a polar ring which gave rise to 22 longitudinal subpellicular microtubules, a cylindrical or cone-shaped conoid within the polar ring and numerous electron dense rhoptries. The number of rhoptries visible in individual tachyzoites varied greatly and was dependent to some extent on the plane of section. A maximum of 24 rhoptries was counted in the anterior end of transversely

sectioned tachyzoites. Rhoptries were not seen posterior to tachyzoite nuclei. In longitudinal sections, rhoptries were elongated, club-shaped structures with narrow, dense necks that extended into the conoid. As many as 14-32 electron-dense bodies were observed primarily posterior to the nucleus. Smaller numbers of these dense bodies were found anterior to the nucleus. Unlike rhoptries, dense bodies were generally round or oval in longitudinal sections. Large numbers of micronemes were seen in the anterior end of tachyzoites and only rarely observed posterior to the nucleus. The micronemes were most often arranged in organized arrays or sheets that were orientated parallel to the pellicular membrane or longitudinal axis of the tachyzoite. The number varied greatly in individual tachyzoites but as many as 60-100 micronemes were counted in longitudinal or oblique sections of selected tachyzoites. In addition, a single micropore, located anterior to the nucleus was seen in some tachyzoites. Parasites multiplied by endodyogeny and many were observed in the process of forming 2 progeny zoites within a single tachyzoite. Rarely, as many as 4 zoites with intact nuclei were seen in division but still attached to each other at the posterior end.

A concerted effort has been focused on the in vitro isolation of Neospora-like protozoal parasites from cattle after studies showed that they were the major diagnosed cause of abortion in California dairy cattle. Histologically, the two bovine fetuses from which isolates were first obtained in 1991 had compatible lesions, including multifocal non-supportive encephalitis, and protozoal tissue cysts like those seen in other natural infections of Neospora-like protozoa in bovine fetuses. The immunological reactivity of tissue cysts in the brains of fetus 66 and fetus 93 was also similar to that seen with many of the cysts in naturally infected fetuses which had strong reactions with N.

caninum and *H. hammondi* antisera and occasional reactions to some of the *T. gondii* antisera.

Isolation of these Neospora-like protozoal parasites from aborted bovine fetuses was difficult because fetuses are generally moderately to severely autolyzed at the time of abortion and protozoal tissue cysts are present in a relatively small proportion of the infected fetuses. Previous ultrastructural studies suggested that most of the protozoal cysts in these fetal tissues were affected by autolysis and were probably non-viable (Barr et al. 1991 Vet. Path. 28:110-16). The two fetuses from which isolates were obtained were in comparatively good postmortem condition. This fact, plus the relatively large number of cysts in these fetuses may have been critical factors in the successful isolation of the protozoal parasites. In addition, the isolation methods were modified. In particular, a longer period of trypsinization in preparing the brain material for cultivation, the overnight incubation of the brain homogenate on the monolayer, and use of the 87-3 bovine trophoblastic cell line for the initial parasite isolation were modifications in these procedures which appeared to be particularly helpful in obtaining the BPA1 and BPA2 isolates. Parasite growth was best maintained in 87-3 and CPAE monolayer cells. This contrasts with *N. caninum* and *Hammondia heydorni* which are reported to grow better in bovine monocyte cells (Speer et al. 1988 Infect. and Immun. 50:566-71).

In comparing these bovine protozoal isolates to those of *T. gondii* and *N. caninum* that have been re-isolated from mouse brains, it was found that the bovine protozoal isolates grew more slowly during isolation and after establishment of continuous growth. Whether this reflects a difference in the virulence of the organisms or a difference in adaptation to culture remains to be determined.

By light microscopy, tachyzoites of the bovine isolates were morphologically similar to in vitro tachyzoites of *T. gondii* and *N. caninum*. Cultivated tachyzoites of the bovine isolates had similar immunohistochemical reactions to tachyzoites of *N. caninum*, reacting strongly with *N. caninum* anti-serum and weakly to serum Tg1 which was produced by immunization of a rabbit with tachyzoite lysates of *T. gondii*. These antigenic reactions were distinctly different from those seen with culture-derived tachyzoites of *T. gondii*. Differences in antigenic reactivity of the cultivated BPA1 and BPA2 tachyzoites, as compared to those of tissue cysts in the source fetuses, could be explained by the stage-specific antigen expression of the different parasites and variations in the methods used to produce the antisera (i.e. immunization with cysts, oocysts or tachyzoite lysates). For example, tissue cyst wall antigens that reacted with antiserum to *H. hammondi* appeared to be lacking on tachyzoites of the bovine isolates in vitro. Unfortunately, a direct comparison of different parasite stages was not always possible since tachyzoites were not identified in the brains of the two bovine fetuses and true cysts have not been observed in the BPA1 or BPA2 cultures. Similarly, *N. caninum* tissue cysts and culture-derived *H. hammondi* tachyzoites were not available for comparison. Differences in antigenic expression may also be affected by host-specific factors. To evaluate this possibility, efforts are under way to obtain material from cattle, dogs, rats, cats and mice that have been experimentally infected with *N. caninum* or the bovine isolates so as to make a direct comparison of antigenic reactivity of the parasites in the same host species.

Thus far, characterization of the in vitro isolates from the two aborted bovine fetuses has shown that these parasites are antigenically and/or ultra-structurally distinct from *T. gondii*, *H. hammondi*, *S.*

cruzi, *Besnoitia* spp. and *Frenkelia* spp. These isolates most closely resemble *N. caninum* parasites which have been most extensively studied in the USA and Scandinavia. The similarity between these parasites indicates that the BPA1 and BPA2 isolates belong to the genus *Neospora*. At present little is known about the life-cycle, including the definitive host, of these *Neospora* parasites in dogs or cattle. A better understanding of the biology of these parasites is essential to determine their taxonomic relationship to each other and to other apicomplexan parasites.

Example 2

This example describes an indirect fluorescent antibody (IFA) test for the detection of parasite-specific antibody responses in cattle that were naturally or experimentally infected with *Neospora* parasites. The methods used here are generally as described in Conrad et al. 1993 *J. Vet. Diagn. Invest.* 5:572-578 (1993).

20 MATERIALS AND METHODS

Parasites and antigen slide preparation

Antigen slides were prepared using tachyzoites of the BPA1 described above. Culture media consisted of Dulbecco's Minimum Essential Medium (DMEM) supplemented with 10% (v/v) heat-inactivated adult equine serum, 2 mM L-glutamine, 50 U/ml penicillin and 50 ug/ml streptomycin (DMEM-HS). Tachyzoites of *Toxoplasma gondii* (RH isolate; kindly provided by Dr. J. Boothroyd) were obtained from CPAE monolayer cultures grown in the same medium except that 10% (v/v) heat-inactivated fetal bovine serum was used instead of equine serum. Parasite-infected cultures were maintained in 25 or 75 cm² flasks incubated at 37°C in an atmosphere of 5% CO₂.

Parasites were harvested for antigen preparation when ≥80% of the CPAE cells in the culture flask were infected with clusters of tachyzoites. The infected monolayer was removed from the flask by scraping into the

medium and then passed 3x through a 25 ga needle to disrupt the cells. The suspension was passed through a 5µm filter to remove cellular debris and tachyzoites were pelleted by centrifugation at 1300 xg for 10 min. After removing the supernatant, the pellet was washed twice in sterile phosphate buffered saline pH 7.2 (PBS) and then resuspended in a modified PBS saline (137mM NaCl, 3mM KCl, 3mM $\text{Na}_2\text{C}_6\text{H}_5\text{O}_7 \cdot 2\text{H}_2\text{O}$, 0.4mM $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, 12mM NaHCO_3 , 6mM glucose) to a final concentration of approximately 2,000/µl. Aliquots of 10 µl of tachyzoite suspension were dispensed into each 4 mm well on 12-well heavy teflon coated (HTC) antigen slides. Slides were air-dried at room temperature and stored at -70° C.

Cattle

Test sera was obtained from naturally infected cows that aborted Neospora-infected fetuses, as well as congenitally infected calves. In addition, sera was obtained from two pregnant heifers that were experimentally infected at approximately 120 days gestation with tachyzoites of the BPA1 isolate derived from CPAE cultures. Tachyzoites were obtained from cultures using the procedure described for harvesting tachyzoites for antigen preparation except that the parasites were not washed in PBS and only the inoculum given to each heifer intravenously was filtered to remove cellular debris. After centrifugation, tachyzoites were resuspended in DMEM and administered by inoculation to each heifer so that 3×10^6 tachyzoites were given IV and 5×10^6 were given IM. A control heifer from the same herd and at the same stage of gestation was inoculated with an equivalent amount of uninfected CPAE cell culture material which was prepared and administered using the same procedures as for the infected heifers. Natural or experimental infections were confirmed by identification of Neospora tachyzoites and/or tissue cysts in fetal or calf tissues using an immunoperoxidase test procedure

(Anderson et al. (1991) J Am Vet Med Assoc 198:241-244
and Barr et al. (1991) Vet Path 28:110-116).

For serological comparison with samples from
Neospora-infected cattle, sera were obtained from the
5 following additional sources: 1) cows that aborted
fetuses which did not have lesions or parasites typical
of Neospora infections, 2) weak calves that were
suspected of having Neospora infections, but showed no
lesions or parasites on post-mortem histopathologic
10 examination, 3) 20 heifers that were purchased as
weanlings from a closed beef herd in Todd County,
Nebraska and maintained under strict isolation, on range
conditions at the Agricultural Research Development
Center, University of Nebraska-Lincoln in Mead, Nebraska,
15 4) 20 pregnant heifers that were maintained on pasture in
California and 5) 21 adult beef bulls or cows that were
originally on pasture and then maintained in the same
feedlot as the experimentally infected heifers.

Serum collection and testing

20 Test and control sera were obtained from blood
samples that were collected by venipuncture into
vacutainer tubes without anticoagulant. After storage at
4° C for 2-12 hr, the blood was centrifuged at 500 xg for
10 min and the serum was removed. Sera was stored either
25 at 4° C for <48 hr or frozen at -70° C until tested.

Antigen slides were thawed at room temperature
immediately prior to use. Sera were initially titrated
in 2-fold dilutions from 1:40 to 1:40,960 to determine
the end-point titer. Ten ul of diluted test or control
30 sera were placed in separate wells on the antigen slides.
Slides were incubated at 37° C for 1 hr in a moist
chamber, washed 3x for 5 min each in PBS, and then tapped
gently to remove excess PBS. Fluorescein-labelled
affinity-purified rabbit anti-bovine IgG diluted 1:500 in
35 PBS was added in 10 ul aliquots to each well. Slides
were incubated at 37° C for 30 min, washed 3 times with
PBS for 5 min each wash, tapped to remove excess PBS,

cover-slipped with buffered glycerol (25% [w/v] glycerine in TRIS-HCL: pH 9.0), and examined at 200 magnification using a fluorescence microscope. The end-point titer was the last serum dilutions showing distinct, whole parasite fluorescence.

RESULTS

Natural infections

Sera collected at the time of abortion from 64 cows were tested for serologic reactivity to Neospora antigen (isolate BPA-1) using the IFA test procedure. Aborted fetuses from 55 of these cows had nonsuppurative encephalitis and/or myocarditis which was consistent with a protozoal infection. In addition tachyzoite and/or cyst stages of Neospora were identified by immunohistochemistry in the tissues of these 55 fetuses (Table 3). In the remaining 9 fetuses there was no indication of encephalitis and/or myocarditis and no detectable protozoal parasites. All of the cows that aborted Neospora-infected fetuses had titers of 320 to 5,120 to Neospora antigens (Table 3). Eight of the cows that aborted fetuses with no detectable Neospora parasites had titers ≤ 160 and one had a titer of 320.

TABLE 3: Titers of cow sera collected after abortion of Neospora-infected fetuses to bovine Neospora (BPA1 isolate) antigens.

5	Neospora titer	Number of cows	<u>Neospora tissue stages in fetus</u>		
			cysts	tachys	cysts & tachys
	320	1	0	1	0
10	640	12	2	8	2
	1280	12	2	10	0
15	2560	15	2	11	2
	5120	15	5	7	3

20 Six of the cows that aborted Neospora-infected fetuses were maintained on their 4 dairies of origin so that these cows could be tested repeatedly over a 6 to 12 month period to determine changes in the Neospora titer. Peak titers of 640 to 2,560 were apparent within the first 20 days after abortion in all of the cows (Figs. 1 & 2). Subsequently, the titers of 4 of the cows (Fig. 1, cows 9, 970 & 522; Fig. 2, cow 578) decreased to 640, whereas the titers of cow 3 (Fig. 1) and cow 1328 (Fig. 2) dropped to 160 within 150 days post-abortion. Cows 25 578 and 1328 were rebred and became pregnant again within approximately 50 to 70 days of aborting Neospora-infected fetuses. When these cows were approximately 4 to 5 months pregnant their Neospora titers increased to their original peak levels of 1,280 and remained at this level until the cows gave birth to full-term calves (Fig. 2). 35 The calf born to cow 1328 had a pre-colostral titer of 20,480 and twin calves born to cow 578 both had precolostral titers of 10,240 to the bovine Neospora isolate. Upon necropsy at 2 to 6 days of age, these 40 calves showed mild nonsuppurative encephalomyelitis or focal mononuclear cell infiltrates the brain parenchyma. Neospora tissue cysts were seen in association with

inflammatory lesions in all 3 calves. The post-colostral titers of sera taken from each calf prior to euthanasia were the same as their precolostral titers.

Serologic titers were determined for four additional calves that were diagnosed as having congenital *Neospora* infections based on the presence of characteristic cyst stages in the brain and/or spinal cord which reacted immunohistochemically with antisera to the BPA-1 bovine *Neospora* isolate. *Neospora* was isolated from the brains and/or spinal cords of calves 1-3 and the parasites were grown continuously in vitro, using a previously described method for isolation of *Neospora* from aborted bovine fetuses. At necropsy, calves 1 and 2 had *Neospora* titers of 20,480, calf 3 had a titer of 10,240 and calf 4 had a titer of 5,120. Sera collected from the dam of calf 4 at calving had a *Neospora* titer of 2,560. Precolostral calf sera and sera from the dams of calves 1-3 were not available for testing.

The titers observed in the 7 congenitally infected calves with confirmed *Neospora* infections were markedly greater than those obtained with sera from 4 weak 1-5 day old calves which were suspected of having *Neospora* infections, but showed no histopathologic evidence of characteristic lesions or parasites on post-mortem examination. One of these uninfected calves had a titer of 160, while the others had titers <80 to bovine *Neospora* antigens. Whether or not these calves had received colostrum was not known.

Experimental infections

Repeated sera samples taken from the 3 pregnant heifers prior to experimental inoculation on day 43 had titers of <80 to *Neospora* BPA1 antigens. The 2 heifers that were infected with culture-derived tachyzoites of the BPA1 bovine isolate developed *Neospora* titers of 640 by day 9 and 1,280 by day 18 after parasite inoculation (Fig. 3). The heifer that received uninfected cell culture material had titers <80 to *Neospora* antigens

throughout the experiment. She was euthanitized 32 days after inoculation to remove her fetus which was viable, histologically normal and uninfected, with no detectable titer to *Neospora*. Peak titers for both infected heifers were detected 32 days after parasite inoculation, at which time the fetus of heifer 413 was removed by caesarian section. Histologically, the fetus had inflammatory lesions and numerous *Neospora* tachyzoites present in its central nervous system. In addition, *Neospora* tachyzoites were isolated from fetal tissues and grown continuously in cell culture. Sera collected from the fetus had a titer of 640 to *Neospora* antigens. After her fetus was removed, the *Neospora* titer of heifer 413 fluctuated between 1,280 and 5,120, until day 193 post-infection when it dropped to 640 (Fig. 3). Heifer 416 calved 158 days after parasite inoculation at which time she had a *Neospora* titer of 1,280 (Fig. 3). The calf had a precolostral *Neospora* titer of 10,240 which was the same as the sample which was collected 2 days later, after ingestion of the dam's colostrum. Clinically the calf appeared normal except that it had decreased conscious proprioception in all 4 limbs when examined prior to euthanasia at 2 days of age. There were minimal histological lesions, consisting of focal gliosis in the central nervous system, but no parasites were detected in fetal tissues.

Uninfected cattle

Fifty three of the 61 (87%) adult cattle tested which had no history of *Neospora* infection had titers ≤ 80 , and all but one animal had titers ≤ 160 to both *Neospora* and *Toxoplasma* antigens (Table 4). The pastured cattle that were moved and subsequently maintained under feedlot conditions did not have higher serologic titers to tachyzoites of bovine *Neospora* or *Toxoplasma gondii* than those kept on pasture. End-point titer determinations of all samples from infected or uninfected cattle were always based on whole tachyzoite

fluorescence. However, in testing the apparently uninfected animals, sera samples from 3 of the cows and 7 of the bulls that were housed in the UCD feedlot had parasite fluorescence which was restricted to the apical
5 end of the parasite. This reaction was particularly marked with the 7 sera from bulls which had apical fluorescence titers of 160 to 320 to both *Neospora* and *Toxoplasma*, while the whole parasite fluorescence titer was ≤ 80 .

TABLE 4: Titers of sera from cattle with no evidence of Neospora infection to bovine Neospora (BPA1 isolate) and Toxoplasma gondii.

5	Cattle Number	Location	Neo	Number	Toxo	
			titer	positive	titer	
			positive			
10	20 heifers	Nebraska pasture	≤80 160 320	16 3 1	≤80 160	14 6
15	20 heifers	California pasture	≤80 160	17 3	≤80 160 ND	13 3 4
20	9 cows	UCD feedlot	<80	9	≤80 160	7 2
	12 bulls	UCD feedlot	≤80 160	11 1	≤80 160	10 2
25						

Figure 4 shows the serologic titers of the 61 uninfected adult cattle plus the 9 cows that aborted fetuses without evidence of Neospora infections compared to the titers of Neospora-infected cows at the time of abortion or calving. Although a majority of the infected cattle had titers ≥1280 to Neospora and most of the cattle that had no evidence of infection had titers ≤80, there was some overlap between these groups in the 160 to 640 titer range (Fig. 4).

Example 3

This example describes isolation of DNA encoding nss-rRNA (sequence shown in Figure 5).

40 Parasites

Bovine Neospora isolates BPA-1 BPA-2, BPA-3, BPA-4, BPA-5 were used for DNA isolation. Parasites were harvested for DNA preparation when > 80% of the CPAE cells were infected with large clusters of tachyzoites. The infected monolayer was removed from the flask by scrapping. The tachyzoites in tissue culture media were

pelleted by centrifugation at room temperature, 1300Xg for 10 minutes. The supernatant was removed and the pellet was resuspended in 10 mls sterile physiologically buffered saline (PBS: pH 7.4), passed through a 25 gauge
5 needle three times to disrupt the CPAE cells, and then filtered through a 5 um disc filter (Gelman Sciences, Acrodisc) to remove cellular debris. The filtered material was pelleted at 1300Xg for 10 minutes and washed in PBS (pH 7.4). The supernatant was removed and the
10 tachyzoite pellet was stored at -70°C until used. Uninfected CPAE monolayer cells were processed by the same procedure and used as controls.

Two methods were used to prepare the DNA from the tachyzoites and control CPAE cells. Initially, DNA
15 was prepared as follows. Briefly, the parasite or control cell pellets were suspended in 1.0 ml STE with 0.5% SDS treated with proteinase K (100µg/ml) and RNAase (100µg/ml) then extracted twice with phenol, once with phenol-chloroform-isoamyl alcohol, and once with
20 chloroform-isoamyl alcohol. DNA was subsequently precipitated with ethanol, dried and resuspended in TE buffer. Other DNA samples were prepared with the Isoquick DNA Extraction kit (Microprobe, Corp., Garden Grove, Calif.) following manufacturer's directions.

25 DNA preparations were electrophoresed in 0.8% (w/v) agarose (FMC) Bioproducts) in 0.5 M Tris/borate/EDTA (TBE) buffer (89 mM Tris, 89mM boric acid, 2mM EDTA) gels stained with ethidium-bromide (0.5 ug/ml) and examined under by ultraviolet (UV) light.

30 Amplification of rRNA gene sequences:

DNA sequences were amplified by the polymerase chain reaction (PCR) using a programmable thermal cycler (Perkin-Elmer). Reactions were performed in 50 to 100 ul volume samples containing approximately 50-100 ng of DNA
35 template, 50 mM Tris buffer (pH 8.3), 1.0 mM MgCl₂, 200 mM of each of the four deoxynucleoside triphosphates, 0.5 U of Taq polymerase (Promega) and 100 pM of universal

primer A (5'CCG AAT TCG TCG ACA CCT GGT TGA TCC CCG ACG
ACC GTG GTC TGA ACG GGA G 3') (SEQ ID No.:2) and primer
C (5'GGG CCC TAG GTG GCG CCG ACG ACC GTG GTC TGA ACG GGA
G 3') (SEQ ID No.:3). The PCR cycling parameters
5 consisted of a single step at 94°C for 3 minutes followed
by 30 cycles denaturation at 94°C for 1 minute, 1 minute
of annealing at 55, and 2 minutes of extension at 72 with
a final extension step of 7 minutes. The PCR
amplification product was an approximately 550-bp
10 sequence from the 5' end of the nss-rRNA gene. A more
extensive 1.8 kb sequence of the nss-rRNA gene was
amplified from BPA1 DNA using universal primer A and
primer B (5' CCC GGG ATC CAA GCT TGA TCC TTC TGC AGG TTC
ACC TAC 3') (SEQ ID No.:4).

15 These reactions were performed using 50ul
reaction samples that contained 100pM of each primer, 1
mM MgCl₂, 50-100 ng of template DNA, 50 mM Tris buffer
(pH 8.3), 1.0 mM MgCl₂, a 200 mM of each of the four
deoxynucleoside triphosphates (dATP, dCTP, dGTP, and
20 dTTP), and 0.5 U of Taq polymerase (Promega). The
amplification cycles were performed by an initial
denaturation step at 94°C for 3 minutes followed by 45
cycles of denaturation at 94°C for 1 minute, 1 minute of
annealing at 55°C, and 4 minutes of extension at 72°C
25 with the final extension step for 7 minutes. For both
amplification reactions, the stationary CPAE cell DNA was
used as the negative template control. Sterile water was
used for the PCR reaction condition controls. Aliquots
of each PCR product were sized by comparison with DNA
30 standards (Bioventures, Inc. TN) after electrophoresis
through a 3% (w/v) NuSieve, 1% (w/v) SeaKem agarose gel
(FMC BioProducts, Rockland ME) stained with ethidium
bromide and visualized under UV light.

Amplification products from 3 to 5 reactions
35 were pooled prior to purification to reduce the
possibility of any nucleotide misincorporation errors by
the Taq polymerase during the elongation step of the

newly synthesized complement strain. The PCR amplification products were purified either by gel electroelution or by spin-columns. Two different spin-columns were used at different times. First, Magic PCR Prep DNA purification System (Promega Corp.) was used following manufacturer's directions. Briefly, the products were electrophoresed through a low temperature melting agarose (Low Melt Agarose, FMC BioProducts,). The DNA was visualized in the gel by ethidium bromide staining and the DNA band was excised into an eppendorf tube. The agarose and DNA were heated (70°C) to melt the agarose. The DNA was separated from the agarose using columns and reagents provided in the kit. Later simpler and less time intensive methods were used by purifying the PCR products using the PCR Select II (5 Primer-3 Prime, Inc) columns which do not require electrophoresis and excision of the product in low melt agarose.

DNA sequencing of the purified PCR products were performed following manufacturer's instructions for the PCR Cycle Sequencing System (BRL ds DNA Cycle Sequencing System). The cycling parameters consisted of a two step program after complete denaturation at 95°C for 3 minutes. The first program step amplified DNA using 20 cycles that included 30 seconds for denaturation, 30 seconds for annealing, and 1 minute of extension. The second program step alternated between denaturation (95°C) and extension (72°C) only. Initially, the universal primers (A, C or B) were used to obtain the first nucleotide sequence data from which internal primers could be constructed and used to amplify internal sequences. All sequencing primers were 5' labelled with adenosine 5' [32 P] triphosphate (Amersham). Reactions were heated to 95°C for 5 minutes prior to loading onto either a 6% (w/v) or 8% (w/v) polyacrylamide, 8 M urea (0.4 mm thick) non-gradient gel using a Model S2 Sequencing Gel Apparatus (GIBCO BRL, Gaithersburg, Md). The sequencing gels were fixed in 10%

acetic acid and 10% methanol to remove the urea prior to transfer of the sequencing products in the gel onto filter paper. The gels were dried using a gel drying apparatus (Biorad Gel Drier, X) for 1 to 2 hours at 70-80°C. The membrane filters were autoradiographed using Kodak X-OMAT X-ray film.

DNA Sequence analysis:

DNA sequences were constructed from at least 3 separate reactions to ensure the accuracy of the nucleotide sequence data obtained. Autoradiographs of the sequencing products were read using Hibio MacDNASIS DNA and Protein Sequence Analysis System (Hitachi Software Engineering Co.). This program and the GCG programs (SEQED, Fragment Assembly, Lineup, and Pretty) (University of Wisconsin Genetics Computer Group) on a VMS system facilitated the construction of the DNA sequences.

Example 4

The DNA prepared in Example 3 was used to design primers and probes for the detection of Neospora. The protocol used was as follows.

Oligonucleotide PCR primers:

- 1) Bovine Neospora Forward Primer
(5'-AAGTATAAGCTTTTATACGGCT-3') (SEQ ID NO.:5)
- 2) Bovine Neospora Reverse Primer
(5'-CACTGCCACGGTAGTCCAATAC-3') (SEQ ID NO.:6)

DNA amplification was carried out in a total volume of 50 μ l. The reaction mixture contained 10 mM Tris-HCl (pH 9.0), 50 mM potassium chloride, 0.1% Triton X-100, 1.0 mM magnesium chloride, 200 mM of each deoxynucleoside triphosphates, 0.42 μ M Bovine Neospora Forward primer and 0.384 μ M Bovine Neospora Reverse primer. After precycle denaturation at 94°C for 4 min to reduce nonspecific amplification, 2.5 U of Taq DNA

polymerase (Promega Corp., Madison, WI) was added and the mixture was overlaid with 50 μ l of mineral oil. Amplification was performed in a DNA Thermal Cycler (Perkin Elmer Cetus Corp., Norwalk, CT) for 31 cycles as follows: denaturation at 94°C for 1 min, annealing at 54°C for 1 min, and extension at 72°C for 2 min. The last cycle was given a prolonged extension period of 7 min. After amplification, 5 μ l of each sample or a BioMarker Low (BioVentures, Inc., Murfreesboro, TN) DNA size standard were mixed with 1 μ l of 6X loading dye and electrophoresed on a 3% Nusieve 3:1 agarose gel (FMC Bioproducts). The gel was stained in a 0.5 μ g/ml ethidium bromide solution for 30 min and observed for the presence of amplification products under ultraviolet illumination.

Oligonucleotide DNA probes:

- 3) BPA/Neospora Internal Probe Sequence
(5'-AGTCAAACGCG-3') (SEQ ID No.:7)
- 20 4) Toxoplasma Internal Probe Sequence
(5'-AAGTCAACGCG-3') (SEQ ID No.:8)

Amplification products were denatured in the gel and transferred to nylon membranes (Hybond-N; Amersham Corp., Arlington Heights, IL) by Southern blotting method. DNA was cross-linked to nylon membrane using a Stratalinker UV crosslinker (Stratagene, La Jolla, CA). Prehybridization, preparation of the labeled internal probe, and hybridization were performed as recommended by the manufacturer for the Enhanced Chemiluminescence 3'-oligolabeling and Detection Systems (Amersham). Labeled internal probe was added to a final concentration of 10 ng/ml of hybridization solution and incubated overnight at 30°C with gentle agitation. After hybridization, the membranes were washed twice for 5 min each at room temperature in 5X SSC and 0.1% (w/v) SDS, and then washed twice for 5 min each at room temperature

in 0.5X SSC and 0.1% (w/v) SDS. Membrane blocking, antibody incubations, signal generation and detection were performed as described by the manufacturer. Membranes were exposed to Kodak X-Omat film for 3-10 min.

5 Results

Using the *Neospora*-specific primers 294 bp PCR products were amplified from DNAs of BPA-1 and *Toxoplasma* (RH and BT isolates). In addition, a 350 bp product was amplified from *Sarcocystis cruzi* DNA. No products were produced with DNAs from various bacteria, CPAE cells, and bovine thymocytes. Only the *Neospora*-specific probe hybridized to the *Neospora* amplification product. Similarly, the *Toxoplasma*-specific hybridized only to the *Toxoplasma* amplification product.

15

Example 5

This example describes experimental infections of pregnant cows with culture-derived *Neospora* tachyzoites.

20 Three cows were inoculated with 8×10^6 tachyzoites of the BPA1 *Neospora* isolate (3×10^6 tachyzoites IV, and 5×10^6 tachyzoites IM). These cows were inoculated at 95 days gestation (Cow #412), 100 days gestation (Cow #416), and 105 days gestation (Cow #413).
25 In each case, a *Neospora* fetal infection was confirmed (Cow #412 expelled an infected mummified fetus; Cow #416 gave birth to a calf that was in utero exposed; and an infected fetus was removed surgically from Cow #416). Two control cows were inoculated with uninfected cell
30 culture and gave birth to uninfected live calves.

These cows were kept and rebred without any intervention. All three experimental cows gave birth to seronegative, clinically normal calves (not all post mortem tissues examined to date).

35 The cows were kept and rebred once again. The previously infected cows (#s 412, 416, 413 were then rechallenged by giving them the same inoculum (8×10^6

tachyzoites, divided and given IV and IM) at 89, 83, and 83 days gestation, respectively. Control cows were rebred and observed. Two infected cows (#s 413 and 416) gave birth to live calves which were clinically normal and seronegative to Neospora antigens. The third cow (#412) aborted 27 days post inoculation. The fetus was recovered. Although mild lesions suggestive of Neospora infection were found, Neospora infection, to date, has not been confirmed (formalin-fixed paraffin embedded tissues negative by immunohistochemistry). The cow was rebred and resorbed its fetus. She was rebred again and she aborted once again at 97 days gestation. This second fetus was not recovered. Thus far histopathologic examination of the tissues from the two clinically normal calves indicates that they were not transplacentally infected with Neospora parasites.

This is the first experiment to show that cattle can be protected against Neospora abortion by immunization with culture-derived tachyzoites of the BPA-1 Neospora isolate.

The above examples are provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: THE REGENTS OF THE UNIVERSITY
OF CALIFORNIA

(ii) TITLE OF INVENTION: Bovine Neospora Isolates and Their Uses

(iii) NUMBER OF SEQUENCES: 8

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Robbins, Berliner & Carson
(B) STREET: 201 N. Figueroa Street, 5th Floor
(C) CITY: Los Angeles
(D) STATE: California
(E) COUNTRY: US
(F) ZIP: 90012-2628

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(vii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Berliner, Robert
(B) REGISTRATION NUMBER: 20,121
(C) REFERENCE/DOCKET NUMBER: 5555-309

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (213) 977-1001
(B) TELEFAX: (213) 977-1003

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1747 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AGTCATATGC TTGTCTTAAA GATTAAGCCA TGCATGTCTA AGTATAAGCT TTTATACGGC	60
TAAACTGCCA ATGGCTCATT AAAACAGTTA TAGTTTATTT GATGGTCTTT ACTACATGGA	120
TAACCGTGGT AATTCTATGG CTAATACATG CGCACATGCC TCTTCCTCTG GAAGGGCAGT	180
GTTTATTAGA TACAGAACCA ACCCACCTTC CGGTGGTCTT CGGGTGATTC ATAGTAACCG	240
AACGGATCGC GTTTGACTTC GGCTGCGAC GGATCATTCA AGTTTCTGAC CTATCAGCTT	300
TCGACGGTAC TGTATTGGAC TACCGTGGCA GTGACGGGTA ACGGGGAATT AGGGTTCGAT	360
TCCGGAGAGG GAGCCTGAGA AACGGCTACC ACATCTAAGG AAGGCAGCAG GCGCGCAAAT	420
TACCCAATCC TGATTCAGGG AGGTAGTGAC AAGAAATAAC AACACTGGAA ATTTCAATTC	480
TAGTGATTGG AATGATAGGA ATCCAAACCC CTTTCAGAGT AACAAATTGA GGGCAAGTCT	540
GGTGCCAGCA GCCGCGGTAA TTCCAGCTCC AATAGCGTAT ATTAAAGTTG TTGCAGTTAA	600
AAAGCTCGTA GTTGGATTTC TGCTGGAAGC AGCCAGTCCG CCCTCAGGGG TGTGCACTTG	660

55

GTGAATTCTA GCATCCTTCT GGATTCTTTC AACTTCATT GTGTGGAGTT TTTCCAGGA 720
CTTTTACTTT GAGAAAATTA GAGTGTTC AAGCAGGCTTG TCGCCTTGAA TACTGCAGCA 780
TGGAATAATA AGATAGGATT TCGGCCCTAT TTTGTTGGTT TCTAGGACTG AAGTAATGAT 840
TAATAGGGAC GGTGGGGGC ATTCTGATTT AACTGTCAGA GGTGAAATTC TTAGATTTGT 900
TAAAGACGAA CTAAGCGAA AGCATTGTC AAAGATGTTT TCATTAATCA AGAACGAAAG 960
TTAGGGGCTC GAAGACGATC AGATACCGTC GTAGTCTTAA CCATAAACTA TGCCGACTAG 1020
AGATAGGAAA ACGTCATGCT TGACTTCTCC TGCACCTTAT GAGAAATCAA AGTCTTTGGG 1080
TTCTGGGGGG AGTATGGTCG CAAGGCTGAA ACTTAAAGGA ATTGACGGAA GGGCACCACC 1140
AGGCGTGGAG CCTGCGGCTT AATTGACTC AACACGGGGA AACTCACCAG GTCCAGACAT 1200
AGGAAGGATT GACAGATTGA TAGCTCTTTC TTGATTCTAT GGGTGGTGGT GCATGGCCGT 1260
TCTTAGTTGG TGGAGTGATT TGTCTGGTTA ATTCCGTTAA CGAACGAGAC CTTAACCTGC 1320
TAAATAGGAT CAGGAAGTTC GTGTTCTTGT ATCACTTCTT AGAGGGACTT TGGGTGTCTA 1380
ACGCAAGGAA GTTTGAGGCA ATAACAGGTC TGTGATGCCC TTAGATGTTT TGGGCTGCAC 1440
GCGCGCTACA CTGATGCATC CAACGAGTTT ATAACCTTGG CCGATAGGTC TAGGTAATCT 1500
TGTGAGTATG CATCGTGATG GGGATAGATT ATTGCAATTA TTAATCTTCA ACGAGGAATG 1560
CCTAGTAGGC GCAAGTCAGC AGCTTGCGCC GATTACGTC CTGCCCTTTG TACACACCGC 1620
CCGTCGCTCC TACCGATTGA GTGTTCCGGT GAATTATTCG GACCGTTTTG TGGCGCGTTC 1680
GTGCCCCGAA TGGGAAGTTT TGTGAACCTT AACACTTAGA GGAAGGAGAA GTCGTAACAA 1740
GGTTTCC 1747

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 52 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CCGAATTCGT CGACACCTGG TTGATCCCCG ACGACCGTGG TCTGAACGGG AG 52

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGGCCCTAGG TGGCGCCGAC GACCGTGGTC TGAACGGGAG 40

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCCCGGATCC AAGCTTGATC CTTCTGCAGG TTCACCTAC

39

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AAGTATAAGC TTTTATACGG CT

22

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CACTGCCACG GTAGTCCAAT AC

22

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AGTCAAACGC G

11

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

AAGTCAAACGC G

11

WHAT IS CLAIMED IS:

1. A biologically pure culture of bovine *Neospora*.
5
2. The culture of claim 1, which is selected from the group consisting of ATCC Accession No. _____ (BPA1), and ATCC Accession No. _____ (BPA6).
- 10 3. A method of detecting the presence of antibodies specifically immunoreactive with a bovine *Neospora* antigen in a biological sample, the method comprising contacting the sample with the *Neospora* antigen, thereby forming a antigen/antibody complex, and
15 detecting the presence or absence of the complex.
4. The method of claim 3, wherein the *Neospora* antigen is an isolated *Neospora* tachyzoite.
- 20 5. The method of claim 4, wherein the isolated *Neospora* tachyzoite is BPA1 or BPA6.
6. The method of claim 3, wherein the biological sample is bovine serum.
25
7. The method of claim 3, wherein the antigen is a protein isolated from a *Neospora* culture.
8. The method of claim 7, the isolated
30 protein has a molecular weight of 106 kD or 19 kD.
9. The method of claim 3, wherein the antigen is immobilized on a solid surface.
- 35 10. The method of claim 3, wherein the complex is detected using a labeled anti-bovine antibody.

11. The method of claim 10, wherein the anti-bovine antibody is fluorescently labeled.

12. A method of detecting the presence of
5 *Neospora* in a biological sample, the method comprising contacting the sample with an antibody specifically immunoreactive with a *Neospora* antigen, thereby forming a antigen/antibody complex, and detecting the presence or absence of the complex.

10

13. The method of claim 12, wherein the antibody is a monoclonal antibody.

14. The method of claim 12, wherein the
15 antibody is immobilized on a solid surface.

15. The method of claim 12, wherein the complex is detected using a second labeled antibody.

20

16. The method of claim 15, wherein the second labeled antibody is specifically reactive with a second *Neospora* antigen.

17. The method of claim 15, wherein the second
25 labeled antibody is specifically reactive with the antibody specifically reactive with the *Neospora* antigen.

18. The method of claim 12, wherein the biological sample is bovine neurological tissue.

30

19. A method of detecting the presence of *Neospora*-specific nucleic acids in a biological sample, the method comprising contacting the sample with a oligonucleotide probe which specifically hybridizes with
35 a target *Neospora*-specific polynucleotide sequence, thereby forming a hybridization complex, and detecting the presence or absence of the complex.

20. The method of claim 19, further comprising amplifying the target *Neospora*-specific polynucleotide sequence.

5 21. The method of claim 19, wherein the target *Neospora*-specific polynucleotide sequence is immobilized on a solid surface.

10 22. The method of claim 19, wherein the oligonucleotide probe is labelled.

23. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and an immunogenically effective amount of a bovine *Neospora* 15 antigen.

24. The composition of claim 23, wherein the bovine *Neospora* antigen is an isolated bovine *Neospora* polypeptide.

20 25. The composition of claim 24, wherein the polypeptide is recombinantly produced.

26. The composition of claim 23, wherein the 25 bovine *Neospora* antigen is expressed by a recombinant virus.

27. A method for protecting a bovine animal from infection by bovine *Neospora*, the method comprising 30 the administration of a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an immunogenically effective amount of a bovine *Neospora* antigen.

35 28. The method of claim 27, wherein the bovine animal is a cow or heifer.

29. The method of claim 27, wherein the pharmaceutical composition is administered when the bovine animal is breeding.

5 30. The method of claim 27, wherein the pharmaceutical composition is administered parenterally.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/03174

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 39/002; C12N 1/10

US CL :424/269.1, 93.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/269.1, 93.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, STN
search terms: neospora

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Parasitology, Volume 106, issued 1993, P.A, Conrad et al., "In vitro isolation and characterization of a <i>Neospora</i> sp. from aborted bovine fetuses", pages 239-249, see the entire document.	1-2

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

A

document member of the same patent family

Date of the actual completion of the international search

23 JUNE 1995

Date of mailing of the international search report

29 JUN 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

JEAN C. WITZ

Kelham Tazson

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/03174

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claims 1-2, drawn to a bovine Neospora culture.

Group II, claims 3-18, drawn to a method of detecting a bovine Neospora antigen and antibodies in serum.

Group III, claims 19-22, drawn to a method of detecting bovine Neospora nucleic acids in a sample.

Group IV, claims 23-30, drawn to a pharmaceutical composition comprising a bovine Neospora antigen and a method of immunizing a bovine against bovine Neospora.

The inventions listed as Groups I-IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The inventions of Groups I-IV lack the same or corresponding special technical features in that the inventions have different functions and modes of operations. Further, the antigen may be sequenced and does not require the isolate.